



Charles A. White, DDS, MS, MSD

Jonathan A. Blansett, DDS, MS

Periodontics & Dental Implants

www.drcawhite.com

Phone (479) 631-6074

201 N 34th Street • Rogers, AR 72756

Fax (479) 936-8660

POST-OP INSTRUCTIONS RIDGE AUGMENTATION PROCEDURES

PRESCRIPTIONS: Have all prescriptions filled prior to surgery. Take your medications only as directed. Regular medication prescribed by your physician should be taken as usual, unless you are told to do otherwise. Take PAIN MEDICATION before you begin to feel discomfort, with fruit juice or a full glass of water. Never take pain medication on an empty stomach. Do not use aspirin or products containing aspirin the week prior to or following surgery. If ANTIBIOTICS were prescribed, take them as directed until all have been taken. Patients taking birth control pills should be aware that ANTIBIOTICS could alter the effectiveness of the birth control pill. PERIOGUARD oral rinse or LISTERINE should be used twice a day starting the day after surgery to kill bacteria around the surgical area. PHENERGAN may be prescribed. This is to be used only if you experience nausea. If ATIVAN or any sedative is prescribed YOU MUST HAVE SOMEONE DRIVE YOU FOR THE APPOINTMENT. Refer to “Medications for Periodontal Surgery” sheet for YOUR prescribed or recommended medications.

SWELLING: After 24 hours, a heating pad or moist heat may be used to decrease swelling. When using heat, apply 20 to 30 minutes of each hour. It is also beneficial to take the anti-inflammatory medication IBUPROFEN or ALEVE and to sleep with your head elevated to reduce swelling. If swelling persists after several days and is not resolving, contact Dr. White or Dr. Blansett for instructions.

BLEEDING: Some bleeding is normal during the first day following surgery. Cover your pillow with a towel to prevent staining. If you have a surgical stent, wear it until you return for the first postoperative visit. Purple blood clots sometimes form along the edges of the stent and may bleed for prolonged periods of time. Simply remove the blood clots with moist gauze and apply pressure to the stent to stop bleeding. SPECIAL NOTE: AVOID PEROXIDE, ALCOHOL, CARBONATED BEVERAGES, DRINKING STRAWS, AND SMOKING for at least 24 hours. These things, as well as heat, can contribute to bleeding problems.

NUTRITION: Adequate nutrition is essential for normal healing. The first 24 hours following surgery your diet should be restricted to COLD LIQUIDS. Dietary supplements, such as Instant Breakfast or Ensure, are good. Gradually progress to eating soft foods, which are easily chewed and swallowed. Chew your foods using teeth far removed from the graft sites so that the grafts are not disturbed. Do not skip meals. If you take nourishment regularly, you feel better, gain strength, have less discomfort, and heal faster. Please eat a normal meal before surgery.

HEALING: Ridge augmentations usually exhibit some swelling during the first 2-3 days. By day 4 or 5, the swelling should be resolving.

SUTURE: A non-resorbable suture will be used that will stay in place until healing is complete, usually for 3-4 weeks. If your sutures become loose, please contact Dr. White or Dr. Blansett immediately.

INSTRUCTIONS CONTINUED ON REVERSE

REST: Plan to rest the remainder of the day of surgery and the following day. After that time, you should be able to return to your regular schedule, EXCEPT such activities as jogging, aerobic exercise, and other strenuous activities. These should be discontinued for the week following surgery. Use good judgement when returning to more strenuous activities.

HEMOCARE: Don't brush or floss near the ridge augmentation for the first week. After 1 week, you may be instructed to brush gently near the augmentation with a very soft toothbrush. You can brush and floss you other teeth as usually but don't rinse you mouth vigorously. Rinse with Periogard or Listerine to keep bacterial counts down in the area of the ridge augmentation.

PAIN: Following surgery, pain is usually mild to moderate in severity and usually persists for 24 to 36 hours. Your discomfort should gradually diminish after the first day. IF YOU HAVE SEVERE PAIN AFTER 24 HOURS, CONTACT DR. WHITE OR DR. BLANSETT. Please make sure that you have no plans to fly for at least one week following surgery. One additional week would be best. Flying could retard healing and cause unnecessary pain and bleeding due to pressure. Automobile trips are fine as long as the surgery is not extensive. Keep in mind that if you leave town following surgery you may not be able to be seen as quickly by Dr. White or Dr. Blansett if there is a problem. We want the opportunity to see you if there are any problems following surgery.

DENTAL IMPLANT PATIENTS: Ridge augmentations requiring bone grafting for dental implants may require the placement of screws and/or tacks that will need to be removed. The screws and tacks will be removed during the placement of dental implants, usually after 4-5 months healing.

QUESTIONS/PROBLEMS: : It is our sincere desire to have you as comfortable as possible during the healing period. If you have any questions or concerns, please don't hesitate to call. Office phone 479-631-6074. Dr. White's cell phone: 479-644-1087. Dr. Blansett's cell phone: 479-282-9448. (Phone calls only – NO texts Please.)